

THE EFFECT OF PREPREGNANCY BMI AND PARITY ON THE DURATION OF BREASTFEEDING DURING EXCLUSIVE BREASTFEEDING

Regina Vidya Trias Novita¹, Pratiwi Harnussant²

STIK Sint Carolus Jakarta, Indonesia

E-mail: reginanovita04@gmail.com*

KEYWORDS	ABSTRACT
BMI; Exclusive Breastfeeding; Pre-Pregnancy; Parity	Obesity Is a condition in a person who experiences overnutrition, Especially Fat Accumulation, Which cause the body to weight more than normal body weight, Indicated with A Body Mass Index (BMI) of more than 29.9 kg/M2 several International Pieces Of Research State That Obese Mothers Fail To Breastfeed Exclusively. Aside From Maternal BMI, Maternal Characteristics Also Support Exclusive Breastfeeding. Maternal Characteristic Factors Include Age, Parity, And Mother's Education. This Study Aims To Determine The Correlation Between Prepregnancy BMI And Maternal Characteristic Factors With The Duration Of Breastfeeding During Exclusive Breastfeeding. This Research Is Quantitative Research Using A Cross-Sectional Descriptive-Correlational Method With A Retrospective Approach. The Number Of Samples In This Research Was 300 Mothers Who Met The Inclusion Criteria With The Purposive Sampling Technique. Data were collected and processed statistically using the Chi-Square Test and T-test to determine factors related to breastfeeding duration, and the linear regression test with the backward method was used to determine the most influential factors in breastfeeding. The Chi-Square Test Results Showed That All Variables Were Related To The Duration Of Breastfeeding With A P-value <0.05. The Linear Regression Test Showed That The Factors That Had A Significant Effect On The Duration Of Breastfeeding Were Prepregnancy BMI And Parity With A Significance Of 0.00; judging From The Coefficient B Factor Value Of 0.285, Parity Had A More Dominant Effect.

INTRODUCTION

Breastfeeding Is Recognized Globally As A Vital Public Health Priority. According To The World Health Organization (WHO), Exclusive Breastfeeding (EBF) For The First Six Months Of Life Is Critical For Infant Health, Providing Optimal Nutrition And Protecting Against Infectious Diseases, Including Respiratory Infections And Diarrhea, Which Are Leading Causes Of Infant Mortality (WHO, 2020). EBF Is Also Known To Contribute To The Physical And Cognitive Development Of Infants, Reducing The Risk Of Chronic Conditions Such As Obesity, Diabetes, And Cardiovascular Diseases Later In Life (Rollins, 2016). Despite These Known Benefits, Global Rates Of Exclusive Breastfeeding Remain Below The WHO's Recommended Target Of 50% By 2025 (WHO, 2020). Factors Influencing The Success And Duration Of Breastfeeding Are Thus Critical Areas Of Research To Achieve Global Health Goals. Several Maternal Factors, Such As Prepregnancy Body Mass Index (BMI) And Parity (Number Of Previous Pregnancies), Have Been Identified As Influential Determinants Of Breastfeeding Duration.

Prepregnancy BMI, An Indicator Of Nutritional Status And Health Before Conception, Can Affect Lactation Success. Studies Suggest That Both Underweight And Obese Women May Face Challenges In Initiating And Sustaining Breastfeeding Due To Physiological Or Hormonal Imbalances (Haugen, 2018). Parity, On The Other Hand, May Impact Breastfeeding Behavior Due To Maternal Experience And Confidence Gained From Previous Pregnancies, Influencing The Duration And Exclusivity Of Breastfeeding (Da Silva, 2020). Understanding How These Factors Affect Exclusive Breastfeeding Duration Is Critical In Addressing Global Breastfeeding Challenges. Research on prepregnancy BMI and breastfeeding yielded mixed results. Some Studies Have Found That Women With Higher BMI Are Less Likely To Initiate Breastfeeding And Tend To Have Shorter Breastfeeding Duration (Wang, 2020). The Underlying Mechanisms May Involve Delayed Onset Of Lactogenesis, Lower Prolactin Responses, And Reduced Milk Supply Among Obese Women (Rasmussen, 2018). However, Other Research Indicates That Overweight Or Obese Women Can Successfully Breastfeed If They Receive Adequate Support And Education On Breastfeeding Practices (Zhu, 2021). Consequently, The Role Of Prepregnancy BMI On Breastfeeding Duration Remains A Topic Of Debate, Requiring Further Investigation To Understand The Complex Relationship.

Obesity Is A Condition in a Person Who Experiences Overnutrition, Especially Fat Accumulation, Which Causes The Body To Weigh More Than Normal Body Weight, Indicated With A Body Mass Index (BMI) Of More Than 29.9 Kg/M², According To The Institute Of Medicine (IOM). Lambert Et Al. (2018) states that 55% Of Women Of Reproductive Age In America Are Obese, And The Rate Increases by 0.5% Points Per Year. Wojcicki (2011) States That More Reproductive-Age Women Are Obese Than Men In The United States, Which Has Become A Global Issue In Recent Years. A Similar Pattern Is Also Found In Indonesia, Where The Prevalence Of Women Of Reproductive Age With Obesity Is 29.3%, Which Is More Than That Of Obese Men, which is 14.5% (Risksedas, 2018). Nobesity In Women Can Pose Several Maternal Risks For Both Mothers And Their Babies. In Pregnant Women, It Can Increase The Occurrence Of Gestational Diabetes Mellitus, Hypertension In Pregnancy, Pre-Eclampsia, Macrosomia, And Labor Complications. The Risk of a Fetus From An Obese Mother Can Cause Neonatal Trauma, Such As Head Trauma And Shoulder Dystocia, And In The Long Run, Can Cause Childhood Obesity (Farpour-Lambert et al., 2018).

Oddy et al. (2006) state that Prepregnancy Obesity Is a risk Of Pregnancy And Labor, Which Can Cause Failure Of Exclusive Breastfeeding Indicated With Early Cessation Of Breastfeeding. One Of The Possible Causes Of Early Breastfeeding Cessation Is The Hormonal Factors, Where The Lactogenesis Process, That Is The Process Of Breast Milk Production And Secretion, Is Delayed. Another Factor Is The Difficulty In Breastfeeding Techniques, Such As Positioning And Attachment. Psychological Factors That Affect The Occurrence Of Breastfeeding Failure Are Changes In Body Shape Where Mothers Feel Ashamed To Breastfeed In Public Places (Wojcicki, 2011). According To Rasmussen And Dewey In Oddy et al. (2006), Women With Prepregnancy Obesity Have A Reduced Prolactin Response To Breastfeeding And An Increase In Progesterone, Which Interferes With The Early Lactation Process. In General, Women With Obesity Experience A Decrease In Prolactin Response On The Seventh Day.

According To Marshall Et Al. (2019), It Is Implicated As The Cause Of Delayed Lactogenesis Which Contributes To The Dysfunction Of Lactation Due To Insufficient Breast Milk Production. An Increase In Prepregnancy BMI And Body Fat Percentage In Early Pregnancy Is Linked To A Significant Reduction In Exclusive Breastfeeding At 6-Week Postpartum. Ramji Et al. (2016) state that Obese Women Are Less Likely To Start And Continue Breastfeeding Due To Anatomical Differences, Low Prolactin Response, And Delays In Lactogenesis II. Several Pieces Of Research On The Correlation Between Obesity

And The Duration Of Breastfeeding That Has Been Carried Out Found Other Factors That Influence Exclusive Breastfeeding, Such As Psychological, Sociocultural, Mother's Intention Of Breastfeeding, And Maternal Sociodemographic Factors. Boudet-Berquier Research (2018) states that The Duration Of Breastfeeding Is Influenced By The Social Environment, Especially Social Norms About Breastfeeding And Public Support After Comparing The Differences In The Average Duration In Several Countries. O'Sullivan Et Al. (2015) States That Multiparous Women Who Are Significantly Obese Are Closely Related To Financial Conditions, Mother's Education, Type Of Delivery Methods, And Breastfeeding Knowledge.

Research On The Influence Of These Factors Has Also Been Carried Out In Indonesia. Mabud's Research (2014) States That Parity Is A Factor That Correlates With Exclusive Breastfeeding Based On The Mother's Experience. Yuliawati Et Al. (2018) states that a mother's education influences breastfeeding success, and the higher the mother's education, the better the knowledge. This Is Because Mothers With Higher Education Know That Giving Exclusive Breastfeeding For Their Babies Is The Best Choice, According To The Information That Respondents Received From Health Workers. It Is Also Possible That Maternal Age Is One Factor That Influences The Success Of Exclusive Breastfeeding. The Results Of Conita's Research (2014) State That Mothers of healthy Reproductive Age (20-35 Years) Have A Higher Success Rate Of Breastfeeding Compared To Mothers Under 20 Years Of Age Associated With Mental Maturity And Pregnancy-Related Conditions.

Government Regulation Number 33 The Year 2012 Recommends Breast Milk Be Given To Newborns For Up To Six Months To Meet Their Nutritional Needs, Without Adding And Or Replacing With Other Food Or Drinks (Except Medicine, Vitamins, And Minerals). Breast Milk Is A Living Liquid Containing Colostrum, Rich In Antibodies Because It Contains Protein For The Body's Immune System And A High Amount Of Germ Killers. So It Can Reduce The Risk Of Death In Infants. Yellow Colostrum Is Produced From The First Day To The Third Day. Breast Milk Contains Less Immunoglobulin, Protein, And Lactose Than Colostrum But Has Higher Fat And Calories With a White Color From The Fourth To The Tenth Day. Breast Milk Also Contains Absorbent Substances In The Form Of Separate Enzymes That Will Not Interfere With Enzymes In The Intestines (Peraturan et al. Indonesia, 2012).

The Success Rate Of Exclusive Breastfeeding In Indonesia Is Quite Good, Reaching 67.74%, And The Special Region Of Yogyakarta (DIY Province) Has Reached 77.50% (Risesdas, 2018). However, The Exclusive Breastfeeding Coverage Rate In Yogyakarta City Has The Lowest Percentage Compared To Other Districts In DIY, Which Is 61.1%. The Number Of Exclusive Breastfeeding Coverage In Yogyakarta Has Also Experienced A Decrease From The Previous Year Which Was 66.1%. The Percentage Of Obese Women Of Their Reproductive Age In Yogyakarta Is Quite High, Which Is 66% (Dinkes et al., 2019). Based On The Above Phenomenon, The Researchers Are Interested In Proving The Influence Of Prepregnancy Obesity And Maternal Characteristic Factors On The Success Of Exclusive Breastfeeding.

Parity refers to the number of times a woman has given birth, which has consistently been associated with breastfeeding duration. Multiparous women (Those With Two or More Pregnancies) are more likely to have longer durations of breastfeeding than primiparous women (Those With Only One Pregnancy) (Girard, 2019). The Experience And Confidence Gained From Previous Pregnancies Often Result In Better Breastfeeding Practices, Fewer Perceived Barriers, And Higher Motivation To Continue Breastfeeding. Additionally, Multiparous Women Are Likely To Receive More Social Support From Family And Friends, Further Facilitating Longer Breastfeeding Periods (Li, 2021). Investigating How Parity Influences Exclusive Breastfeeding Duration Is Essential For Developing Targeted Interventions For First-Time Mothers. There Is Growing Evidence That Prepregnancy BMI

And Parity May Interact To Influence Breastfeeding Outcomes. For Instance, Studies Have Shown That Among Multiparous Women, The Negative Effect Of High Prepregnancy BMI On Breastfeeding Duration Is Less Pronounced Than Among Primiparous Women (Kent, 2020). This Suggests That Experience Gained From Previous Breastfeeding Attempts May Mitigate Some Of The Physiological And Psychological Barriers Associated With Higher BMI. Understanding How These Two Factors Jointly Impact Exclusive Breastfeeding Duration Can Help Tailor Breastfeeding Interventions To Different Maternal Profiles. From A Theoretical Standpoint, The Physiological And Psychological Components Of Breastfeeding Are Intertwined. Hormonal Regulation, Particularly The Roles Of Prolactin And Oxytocin, Is Central To Lactation Success. These Hormones Regulate Milk Production And Ejection, But Their Effectiveness Can Be Influenced By Maternal Factors Such As BMI And Psychological Stress (Brown, 2018). Furthermore, the Self-Efficacy Theory posits that a mother's confidence in her ability to breastfeed plays a critical role in determining the duration of breastfeeding (Dennis, 2019).

Parity Enhances Maternal Experience and is Likely To Increase Self-Efficacy, Thereby Promoting Sustained Breastfeeding. Several Studies Have Explored The Effects Of Prepregnancy BMI And Parity On Breastfeeding Outcomes. A Systematic Review (Castilho, 2020) Found That Women With Higher Prepregnancy BMI Had Lower Rates Of Breastfeeding Initiation And Shorter Durations Of Exclusive Breastfeeding. Similarly, A Meta-Analysis By (Torres, 2021) Concluded That Multiparous Women Were More Likely To Breastfeed Exclusively For Six Months Compared To Primiparous Women. However, research on the combined effect of BMI and parity is still limited, and further investigation is warranted. Despite The Growing Body Of Literature On Maternal Factors Influencing Breastfeeding, There Remains A Gap In Understanding How Prepregnancy BMI And Parity Interact To Affect Exclusive Breastfeeding Duration. Previous Studies Have Predominantly Examined These Factors In Isolation Without Considering Their Combined Effect On Lactation Outcomes (Morales, 2021).

This Study Aims To Fill This Research Gap By Investigating The Interaction Between Prepregnancy BMI And Parity In Determining Exclusive Breastfeeding Duration. The Primary Objective Of This Research Is To Assess The Effect Of Prepregnancy BMI And Parity On The Duration Of Exclusive Breastfeeding. Specifically, The Study Aims To Examine The Relationship Between Prepregnancy BMI And Exclusive Breastfeeding Duration, Explore How Parity Influences Breastfeeding Behavior And Outcomes, Investigate The Combined Effect Of Prepregnancy BMI And Parity On The Likelihood And Duration Of Exclusive Breastfeeding, Provide Insights Into How Health Interventions Can Be Tailored To Improve Breastfeeding Outcomes For Women With Varying BMI And Parity Profiles.

METHOD RESEARCH

This Research Used A Quantitative Method With An Analytical Observational Type Of Research Aimed To Get A prediction of A Population And Find Out The Correlation Between Cause And Effect Or Risk Factors With Effects. The Design Used For This Research Was A Retrospective Cross-Sectional Design. The sampling method used was non-probability sampling, and purposive sampling was used. The Inclusion Criteria Were Mothers Who Gave Birth In 2019 And Were Domiciled In Yogyakarta. The Samples Obtained Were 300 Mothers, With The Proportion Of 150 Mothers With Normal BMI As A Control Variable And 150 Mothers With Obese BMI. The Research Instrument Was A Brief Questionnaire Containing The Data On Maternal Weight Before The Last Pregnancy, Mother's Last Education, Age, Parity, Duration Of Breastfeeding, And The Time When Additional Food Was Given To The Babies For The First Time.

This Research Used The Chi-Square Test And T-Test Analysis To Determine The Relationship Between Prepregnancy BMI And Maternal Characteristic Factors With Breastfeeding Duration, With A Significance Level Of $\alpha = 0.05$ And If $T\text{-Count} > T\text{-Table}$. A Linear Regression Test Was Conducted To Determine The Most Influential Variable On The Duration Of Breastfeeding. The Analysis Of This Research Used The SPSS Version 21 Program. Data Collection Was Carried Out In May – July 2021 In All Public Health Centers In The Yogyakarta City Health Office Area. The Researchers Carried Out Data Collection By Conducting Short Online Interviews. In Some Areas Of The Public Health Center, The Researchers Were Assisted By Cadres In Collecting Samples By Distributing Online Questionnaires.

RESULT AND DISCUSSION

Table 1 shows that the research respondents ranged from 17 to 48 years old, with the majority of respondents (81.3%) being of reproductive age (20 – 35 years) and the average age being 29.47. The Majority Of Respondents In This Research Were Primiparous (56.3%) And The Majority Of Mothers Had Higher Education (93.7%). Mother's Prepregnancy BMI Ranged From 17.30 – 43.40 With An Average Value Of 25.68 Where This Number Refers To Obese BMI. Based On The Data Containing The Duration Of Breastfeeding That Has Been Collected, The Majority Of Mothers Managed To Breastfeed Exclusively (64.3%) With A Duration Range Of 0-6 Months. Respondents Breastfed Exclusively For 4.65 Months On Average.

Table 1. The Frequency Distribution Of Respondents' Characteristics In Yogyakarta

Characteristics	N	%	Mean	Min	Max
Age					
< 20 Yo	9	3			
20 – 35 Yo	244	81.3	29.47	17	48
>35 Yo	47	15.7			
Parity					
Primi	169	56.3			
Multi	126	42	1.74	1	5
Grande	5	1.7			
Education					
Low	19	6.3			
High	281	93.7	-	-	-
Prepregnancy BMI					
Normal BMI	150	50	25.68	17.30	43.40
Obese BMI	150	50			
Breastfeeding Duration					
≤ 2 Months	90	30			
4 Months	17	5.7	4.65	0	6
6 Months	193	64.3			
Total	300	100			

Source: Data Primer, 2021

Table 2. The Effect Of Prepregnancy BMI On The Duration Of Breastfeeding During Exclusive Breastfeeding

BMI	Breastfeeding Duration								BMI	T-Value	P-Value
	≤ 2 Months		4		6		Total				
			Months		Months						
	N	%	N	%	N	%	N	%			
Normal	32	21.3	4	2.7	114	76	150	100	21.57	112.5	.000
Obese	58	38.7	13	8.7	79	52.7	150	100	29.80	69.4	.000

Source: Data Primer, 2021

Table 2 Shows The Results Of The Bivariate Analysis Of The Prepregnancy BMI Variable. 76% Of Mothers With Normal BMI Had The Longest Breastfeeding Duration Of 6 Months, And 21.3% Of Mothers With Normal BMI Did Not Succeed In Exclusive Breastfeeding With A Breastfeeding Duration Of ≤ 2 Months. The Average Value Of BMI In Mothers With Normal BMI Is 21.57. The Result Of The Statistical Test On Mothers With Normal BMI Had A Significance Value Of 0.00, And The Analysis Of the T-Value Was $112.5 > 1.968$, Which Means That Normal BMI Corresponds To The Duration Of Breastfeeding During Exclusive Breastfeeding And Has 112.5 Chances For Exclusive Breastfeeding. Data On Mothers With Obese BMI Showed That 58 (38.7%) Mothers Had Breastfeeding Duration Of ≤ 2 Months, 13 (8.7%) Mothers Exclusively Breastfed For Up To 4 Months, And 79 (52.7%) Obese Mothers Managed To Breastfeed For 6 Months. The Average Value Of BMI In Obese Mothers Is 29.80. The Chi-Square Statistical Test Results Showed That Mothers With Obesity Showed A P-Value Of 0.00. From The Analysis Of The T-Value Of $69.3 > 1.968$, It Showed That Obese BMI Corresponds To The Duration Of Breastfeeding With 69.3 Chances Of Exclusive Breastfeeding.

Table 3. The Effect Of Characteristics On The Success Of Exclusive Breastfeeding

		Breastfeeding Duration						Total		T-Value	P-Value
		≤ 2 Months		4 Months		6 Months					
		N	%	N	%	N	%	N	%		
Age		90	30	17	5.7	193	64.3	300	100	89.02	.001
	<20 Year	5	55.6	2	22.2	2	22.2	9	100		
	20-35 Ear	78	32	15	6.1	151	61.9	244	100		
	>35 Year	7	14.9	0	0	40	85.1	47	100		
Education		90	30	17	5.7	193	64.3	300	100	137.49	.004
	Low	0	0	0	0	19	100	19	100		
	High	90	32	17	6	174	61.9	281	100		
Parity		90	30	17	5.7	193	64.3	300	100	47.39	.000
	Primi	74	43.8	13	7.7	82	48.5	169	100		
	Multi	16	12.7	4	3.2	106	84.1	126	100		
	Grande	0	0	0	0	5	100	5	100		

Source: Data Primer, 2021

Table 3 Shows The Characteristics Of The Mothers, While The Age Factor Shows That Mothers Who Are In Reproductive Age (20-35 Years) Have A Success Rate Of Breastfeeding For 6 Months, With The Percentage Of 61.9%, Mothers With Age > 35 Years Also Show A Fairly High Percentage Of Breastfeeding Duration Up To 6 Months That Is Equal To 85.1%. In Mothers Aged < 20 Years, It Was Discovered That 55.6% Did Not Give Exclusive

Breastfeeding With A Breastfeeding Duration Of ≤ 2 Months. The Result Of Statistical Tests Using The Chi-Square Test Found A P-Value Of 0.000; Therefore, The Significance Level Was <0.05 , And The T-Value Is $89.02 > 1.968$, Which Means That Research Shows A Correlation Between Age Characteristics And Breastfeeding Duration. In Terms Of Educational Characteristics, Most Of The Respondents Have A High Level Of Education; from A Total Of 300 Respondents, Only 19 Respondents Have A Low Level Of Education. Mothers With Low Education Were All Successful In Exclusive Breastfeeding (100%), While Mothers With High Levels Of Education Had A Percentage Of 61.9% For Breastfeeding With A Duration Of 6 Months.

The Results Of Statistical Tests Using The T-test showed That The T-value was 66,499 $> 1,968$, And The Chi-Square Test Found A P-value of 0.001. Therefore, It Is Concluded That There Is A Correlation Between A Mother's Education And The Duration Of Breastfeeding. The Results Of the Parity Factor Analysis Showed That Mothers Who Had Given Birth Previously Had A High Success Rate Of Exclusive Breastfeeding. Multigravida Mothers Have A Success Rate Of 84.1%, And All Grand Multigravida Mothers Are Successful In Giving Exclusive Breastfeeding (100%). Mothers Who Gave Birth For The First Time (Primigravida) Showed The Highest Failure Rate In Exclusive Breastfeeding Compared To Other Groups Of Mothers, Which Was 43.8% With A Breastfeeding Duration Of ≤ 2 Months. The Result Of Statistical Tests Using The Chi-Square Test Found A P-Value Of 0.000 <0.05 , And The Result Of The T-Test Value Was $14,784 > 1,968$, Which Means It Can Be Concluded That There Is A Correlation Between Maternal Parity And Duration Of Breastfeeding.

Table 4. The Result Of The Linear Multivariate Regression Test With The Backward Method Between Characteristics And Prepregnancy BMI On The Duration Of Breastfeeding In Exclusive Breastfeeding

	Sig.	Coef (B)	T	95% C.I For EXP (B)	
				Lower	Upper
Prepregnancy BMI	.000	-.222	-4.162	-.594	-.212
Age	.072	.108	1.803	-.022	.497
Education	.022	-.122	-2.298	-.847	-.066
Parity	.000	.285	4.741	.286	.692
Constant	.000		5.485	1.677	3.554

Source: Data Primer, 2021

Table 4 Is The Result Of The Linear Regression Analysis; two Variables Have A Significance Of > 0.05 , Which Are The Educational Variable With A Value Of 0.22 And Age With A Value Of 0.072. This Means That Both Age And Education Do Not Have A Significant Effect On The Duration Of Breastfeeding During Exclusive Breastfeeding; Therefore, Both Variables Are Excluded From The Characteristic Factor. In The Prepregnancy BMI Variable And The Parity Variable Both Have The Same Significance Level Of 0.00. Therefore, The Significance Level Is <0.05 ; it Is Concluded That The Prepregnancy And Parity BMI Variables Have A Strong Significant Effect On The Duration Of Breastfeeding During Exclusive Breastfeeding. Both Variables Have The Same Significance Level, Based On The Value Of The B Coefficient, It Can Be Concluded That Parity Is A More Positive Factor Affecting The Success Of Exclusive Breastfeeding Where The B Coefficient Is 0.285 And The T-value is 4.741, While The B Coefficient On Prepregnancy BMI Is -0.222 With A T Value Of - 4.162 Which Means, The Higher The Value Of A Mother's Prepregnancy BMI, The Shorter The Duration Of Breastfeeding During Exclusive Breastfeeding.

DISCUSSION

Research Data Show That Mothers With Obesity Before Pregnancy Have A Lower Success Rate Of Exclusive Breastfeeding Compared To Mothers With Normal Weight. The Results Of This Research Are Aligned with Infants Of Obese Mothers Have Donath Et Al. (2008) And Boudet-Berquier Et Al. (2018), Which shows that Obese Women Are Inclined Not To Breastfeed Exclusively. The Results Of The Research Conducted By Donath & Amir (2008) Entitled "Maternal Obesity And Initiation And Duration Of Breastfeeding: Data From The Longitudinal Study Of Australian Children" State That Obese Women Stop Breastfeeding In The First Week Of Birth Compared To Women Of Normal Weight And Those Who Manage To Breastfeed For 1 Week Tends To Stop Before 6 Months. Berquier's Research (2018), Entitled "Association Between Maternal Prepregnancy Obesity And Breastfeeding Duration: Data From A Nationwide Prospective Birth Cohort," Also Has The Same Results Where Obese Women Tend To Give Mixed Formula Milk Earlier Than Women With Normal BMI. The Results Of This Research State That 83% Of Obese Mothers Started Giving Additional Food Or A Mixture Of Formula Milk To Their Babies At The Age Of 3 Months. Prepregnancy Obesity Is Thought To Be The Strong Cause Of The Failure Of Exclusive Breastfeeding Compared To Mothers Who Experience Excessive Weight Gain During Pregnancy. Women Who Are Obese Long Before Pregnancy Tend To Have A Disturbance In Insulin Balance, Which Results In Impaired Protein In The Body That Can Affect Breast Development, Which Leads To Obese Women's Breasts Being Inclined To Have Thicker Fat Tissue And Fewer Mammary Glands (Guyton & J.E., 2007).

In Contrast To Bartok Et Al. (2012) And O'Sullivan's (2015) Research Stating That The BMI Of Obese Mothers Was Not An Independent Risk Factor In The Failure Of Exclusive Breastfeeding; the Last Multivariable Factor That Influenced The Success Of Exclusive Breastfeeding The Most Was How Long The Mother Planned To Breastfeed Her Baby And The Low Self-Efficacy Factor. Mothers With Prepregnancy Obesity Also Experience Insulin Resistance, Which Is Known That In Early Childbirth, Milk Production And Secretion Are Regulated By The Endocrine And Require Insulin. The Insulin Resistance Found In Obese Mothers Inhibits Milk Production, Which Causes Delays In The Process Of Lactogenesis II In Obese Mothers (Preusting et al., 2017). Adipose Tissue Or Thick Fat In Obese Women Allows Higher Levels Of Progesterone Because The Storage Of Progesterone Occurs In Fat. It Is Known That High Progesterone Levels Will Hold Or Inhibit Breast Milk Secretion (Marshall et al., 2019). The Delay In The Process Of Lactogenesis II And Disturbances In The Production Of Breast Milk Make Obese Mothers Unable To Meet The Needs Of Their Baby's Milk At The Beginning Of Birth—High Fat Levels in Obese Mothers Increase Leptin Secretion. High Levels Of Leptin Increase The Performance Of Estrogen Receptor Alpha (Era), Thus Causing High Levels Of Estrogen In The Body Of Obese Mothers; Therefore, It Interferes With Or Reduces Breast Milk Production (Chamberlin et al., 2017). Research Done By Oddy (2006) States That Infants Of Obese Mothers May Have High Energy Needs. The Problem Of Low Milk Production And the High Needs For Breastfeeding For Infants Makes The Infants' Satisfaction With Breastfeeding Unfulfilled, Allowing Obese Mothers To Stop Exclusive Breastfeeding Even Though They Have Successfully Breastfed In The Early Weeks Of Birth, And There Is A Possibility That Mothers Will Start Making Mixed Nutrition With Formula Milk.

Based On The Results Of This Research, The Researchers Assume That Prepregnancy Obesity Influences Exclusive Breastfeeding. This Is Likely To Happen Because Mothers With Prepregnancy Obesity Have Different Breast Anatomy Conditions And Hormonal Patterns. This Will Affect Milk Production. 38.7% Of Obese Mothers Stopped Breastfeeding After ≤ 2 Months; this May Be Due To A Lack Of Milk Production In The Early Postnatal Period That

Leads To Mothers Feeling Their Milk Is Not Coming Out And Deciding To Stop Breastfeeding. The Results Of The Data Analysis Show That Mothers With Obesity Have 96 Opportunities To Breastfeed Exclusively. Maternal Characteristics In This Research Probably support this; the Majority Are Of Reproductive Age And Have Higher Education. In Terms Of Age Characteristics, This Research Is In Alignment With The Research Conducted By Conita (2014), Which States That Age Has An Indirect Correlation With Breastfeeding Success After Linking The Age Factor With The Level Of Emotional Maturity Of The Mother In Taking Care Of Her Baby. In Her Research, Respondents Aged < 20 Years Had The Lowest Success Rate Of Breastfeeding. Failure In Exclusive Breastfeeding Possibly Happens Because They Are Not Physically, Psychologically, And Socially Ready. This Can Disrupt The Psychological Balance That Can Affect Breast Milk Production, Resulting In Mothers Being Unable To Care For Their babies optimally. Several Other Studies State That Age Does Not Correlate With Breastfeeding. Ulfah & Nugroho's Research (2020), Which Examines The Correlation Between Age, Occupation, And Mother's Education In Exclusive Breastfeeding, States That Age Does Not Have A Significant Correlation With Exclusive Breastfeeding With A P-Value Of 0.413. The Same Thing Was Also Shown In Fakhidah & Palupi's Research (2018), Which Stated That There Was No Difference In The Proportion Of Breastfeeding For Mothers Aged 20-30 Years, Aged < 20 Years, And Aged > 35 Years.

Arini (2012) States That Maternal Age Greatly Determines Maternal Health Because It Is Related To The Conditions Of Pregnancy, Childbirth, And Postpartum, As Well As How To Breastfeed The Baby. In This Research, The Success Rate Of Exclusive Breastfeeding Was Mostly Found In The Reproductive Age Group. Mothers Of A Healthy Reproductive Age Have More Mature Physical Conditions And Can Manage Emotions; Therefore, They Are More Prepared To Face Pregnancy, Childbirth, And care for their Babies. As Mentioned By Wawan and Dewi (2011), The Older A Person Is, The Better Their Level Of Maturity And Strength To Think And Work Will Be. According To Arini (2012), Older Mothers Whose Age Is At Risk For Giving Birth Experience A Decrease In Hormone Production, Resulting In A Decreased Lactation Process. At This Age, Giving Birth Is Considered A Risk Because It Is Closely Related To Nutritional Anemia, Which Can Affect Breast Milk Production. In This Research, The Success Rate Of Breastfeeding In Older Mothers Is Quite High. This Is Possible Because Of Another Factor: Where The Older The Mother, The More Emotionally Mature The More Knowledge And Experience She Has, And The Higher Her Awareness And Responsibility Will Be; Therefore, It Affects Breastfeeding Patterns (Conita, 2014). Based on the analysis above, the researchers assume that age-characteristic factors are indirectly related to the duration of breastfeeding. Data showed that 80.1% of mothers aged > 35 years had a 6-month duration of breastfeeding; physiological factors in older mothers did not significantly impact The duration of breastfeeding. In Mothers Aged < 20 Years, 55% Of Mothers Stopped Breastfeeding Exclusively When The Baby Was ≤ 2 Months Old; Therefore, It Can Be Concluded That Age Correlates With The Duration Of Breastfeeding Due To The Maturity And Mental Readiness Of The Mother In Taking Care Of Her Baby.

In This Research, Education Is Correlated With The Duration Of Breastfeeding, which is In Alignment With Yuliawati's Research (2018), Which States That a Mother's Education Has A Correlation With Breastfeeding Failure where The Higher The Mother's Education Level, The Knowledge Will Be Better And It Will Be Easier To Accept A New Idea. Cognita (2014) also states the same thing: the higher a person's level of education is, the higher their ability to absorb knowledge; however, a high level of education without adequate understanding of breastfeeding does not guarantee successful breastfeeding. This Is In Contrast To Ulfah and Nugroho's Research (2020), Which Stated That There Was No Correlation Between The Mother's Level Of Education And The Success Of Breastfeeding. (Fakhidah and Palupi's Research (2018) Also Stated Similar Results That There Was No Correlation Between

Education And The Success Of Exclusive Breastfeeding. According To Her Research, Mothers With Low Education Levels Were No Less Proficient Than Mothers With Higher Education In Seeking Information Through Electronic Media And Information From Cadres Or Midwives.

In General, Maternal Education Is Something That Can Support Exclusive Breastfeeding. Mothers With Higher Education Will Have Broader Knowledge Than Mothers With Lower Education. Education Can Also Encourage A Mother To Seek Experiences That She Has Not Previously Had, Which Can Increase Her Knowledge. Improved knowledge is obtained through formal education. Besides That, support from family is also a supporting factor in exclusive breastfeeding (Ulina & Doveriyanti, 2011). In This Research, It Was Discovered That 100% Of Mothers With Low Education Levels (Elementary And Junior High School) Had A 6-month Breastfeeding Duration, While There Were Still 32% Of Mothers With High Education Levels Who Breastfed With a duration Of ≤ 2 Months. The Researchers Assume that Mothers With Low Education Levels Do Not Also Mean Having Less Knowledge. Knowledge Is Not Always Obtained Through Formal Education; Easy Access To Information Can Be Used To Improve Mothers' Knowledge. The data of this research indicate that mothers who have given birth before (multipara and grand multigravida) have a higher success rate of breastfeeding than mothers who have given birth for the first time (primipara). This Is In Alignment With The Research Done By Ngo Et Al. (2019), Which States That The Experience Of Mothers In Breastfeeding Is A Fast And Strong Foundation Of Successful Breastfeeding and higher Self-Efficacy Than Those Who Have Not Had Breastfeeding Experience; Therefore, The Success Rate Of Exclusive Breastfeeding Is Higher Compared To Mothers Who Just Gave Birth The First Time.

In Contrast, To Untari's Research (2017) Stated That Parity Had Nothing To Do With The Duration Of Breastfeeding, As There May Be Sociocultural Differences, Mothers' And Family's Lack Of Knowledge, And Inadequate Health Services. Similar Results Were Also Found In Fauzi's Research (2019), Which Stated That Parity Was Not Related To The Duration Of Breastfeeding. The Family Support Factor Is Thought To Be An Influence On The Mother's Decision To Breastfeed; Therefore, Mothers Who Have Had Previous Breastfeeding Experiences Do Not Necessarily Have A Positive Influence On The Duration Of Breastfeeding. Primiparous mothers have few prolactin receptors, which causes reduced prolactin stimulation that interferes with milk production. Primiparous Mothers Will Experience Problems When It Comes To Caring For Their Babies Because Of The Lack Of Experience. Education And Experience From Others Will Play An Important Role In Primiparous Mothers. If the mother has poor experience and education, it can cause doubts about the implementation of breastfeeding (International et al. Association In Hasiana et al., 2014). Based On The Results Of The Analysis Above, The Researchers Assumed That Parity Is A Characteristic Factor Associated With The Duration Of Breastfeeding, As Seen in the Data In This Research Which Showed That Multiparous Mothers (84.1%) And Grand Multigravida (100%) Had A 6-Month Breastfeeding Duration. Previous Experience Factors Have A Positive Impact On Breastfeeding. This Is Likely Because Mothers Who Have Previous Breastfeeding Experience Have A Higher Level Of Confidence In Caring For Their Babies Compared To Primigravida Mothers Who Have No Experience At All, Which Can Trigger Anxiety Or Doubts When Breastfeeding Their Babies.

Based On The Linear Regression Test, The Prepregnancy BMI And Parity Variables Were The Most Influential Factors in breastfeeding Duration. Still, The Parity Variable Had A Greater Influence With A B Coefficient Of 0.285. The Results Of This Research Are In Alignment With Ervina's Research (2018), Which States That The Parity Factor Is Related To Exclusive Breastfeeding, Which Shows That Primiparous Mothers Have A 3-fold Risk Of Not Giving Exclusive Breastfeeding Compared To Multiparous Mothers. The Results Of The

Interview Concluded That Primiparous Mothers Have No Experience Lack Knowledge, And Have The Assumption That Breastfeeding Can Make The Breasts Saggy. Multiparous Mothers Already Have Experience And Knowledge From Previous Parity; Therefore, They Have A Greater Chance Of Giving Exclusive Breastfeeding (Mabud et al., 2014). This Is In Contrast To Hastuti et al. (2015) And Untari's (2017) Research, Which Stated That Parity Did Not Affect Exclusive Breastfeeding. According To Their Research, Parity Did Not Affect Exclusive Breastfeeding. Still, sociocultural factors may include mothers' and families' lack of knowledge, unsupportive health lines, and family support factors. If The Mother Has Had Experience With Children Before, But There Were No Supporting Factors, It Would Affect The Mother's Decision To Give Exclusive Breastfeeding.

The Obesity Variable Is Also Statistically Proven To Affect Exclusive Breastfeeding. These Results Are Aligned with The Research Done By Berquier Et Al. (2018), Which States That Obese Women Tend To Stop Breastfeeding Before 6 Months. Berquier Divided Obese Mothers Into Two Groups, Namely Obese Primiparous Mothers And Obese Multiparous Mothers. In Obese Primiparous Mothers, The Failure Of Exclusive Breastfeeding Occurs Due To The Low Prolactin Response And The Delay In The Lactogenesis Process, While The Failure Of Exclusive Breastfeeding In The Group Of Obese Multiparous Mothers Depends On The Method Of Delivery And Weight Gain During Pregnancy. Based On The Analysis Above, The Writer Assumes That The Prepregnancy BMI And Parity Are Two Interrelated Variables. Primiparous Mothers Have A Small Number Of Prolactin Receptors And Lack Of Experience Coupled With Obesity Conditions That Allow For Disturbances In Breastfeeding, Such As Hormone Disorders, Imperfect Breast Growth, And Delays In The Process Of Lactogenesis II, Thereby Increasing The Failure Of Exclusive Breastfeeding In Primiparous Mothers With Obesity (Donath et al., 2008; Guyton & J.E., 2007; Hasiana et al., 2014; Ngo et al., 2019). Multiparous Mothers With Obesity Have A Lower Risk Of Exclusive Breastfeeding Failure Due To The Factor Of A Greater Number Of Prolactin Receptors And Having Had Breastfeeding Experience In Previous Childbirths (Ngo et al., 2019). According To (Hastuti et al. (2015), Multiparous Mothers Who Have Previously Breastfed For > 3 Months Will Breastfeed Their Next Children For A Longer Duration.

CONCLUSION

In Conclusion, Prepregnancy BMI And Parity Are The Most Influential Factors On The Duration Of Breastfeeding For Mothers During Exclusive Breastfeeding, where The Higher The BMI Value, The Lower The Probability Of Exclusive Breastfeeding Is, And The Higher The Parity Of The Mother, The Higher The Probability Of Exclusive Breastfeeding Is. These Two Variables Are Interrelated; Primiparous Mothers Who Are Obese Have A Risk Of Failure To Exclusively Breastfeed Compared To Multiparous Mothers Who Are Obese Due To The Factors Of Breastfeeding Experience And Disruption Of Lactation Processes.

REFERENCES

- Arini. (2012). *Seorang Ibu Harus Menyusui*. Flashbooks.
- Bartok, C. J., Schaefer, E. W., Beiler, J. S., & Paul, I. M. (2012). Role Of Body Mass Index And Gestational Weight Gain In Breastfeeding Outcomes. *Breastfeeding Medicine*, 7(6), 448–456. <https://doi.org/10.1089/Bfm.2011.0127>
- Boudet-Berquier, J., Salanave, B., Desenclos, J. C., & Castetbon, K. (2018). Association Between Maternal Prepregnancy Obesity And Breastfeeding Duration: Data From A Nationwide Prospective Birth Cohort. *Maternal And Child Nutrition*, 14(2). <https://doi.org/10.1111/Mcn.12507>

- Brown, A., & J. S. (2018). Impact Of BMI And Psychological Factors On Breastfeeding Duration: A Biopsychosocial Model. *Journal Of Reproductive And Infant Psychology*, pp. 132–144.
- Castilho, E. M., Et Al. (2020). Prepregnancy BMI And Breastfeeding Duration: A Systematic Review And Meta-Analysis. *Obesity Reviews*.
- Chamberlin, T., D'Amato, J. V., & Arendt, L. M. (2017). Obesity Reversibly Depletes The Basal Cell Population And Enhances Mammary Epithelial Cell Estrogen Receptor Alpha Expression And Progenitor Activity. *Breast Cancer Research*, 19(1). <https://doi.org/10.1186/S13058-017-0921-7>
- Conita, Destia. A. (2014). Perbedaan Pertumbuhan Bayi Usia 3-6 Bulan Yang Diberi Asi Eksklusif Dan Yang Tidak Diberi Asi Eksklusif Di Puskesmas Gang Sehatkecamatan Pontianak Selatandestia Ayu Conitai 111 09 025program Studi Pendidikan Dokterfakultas Kedokteran Universitas Tanjungpurapontianak 2014. *Jurnal Mahasiswa PSPD FK Universitas Tanjungpura*, 1(1). <https://jurnal.untan.ac.id/index.php/jfk/article/download/5495/5669>
- Da Silva, V. F., Et Al. (2020). Parity And Breastfeeding Duration: A Systematic Review And Meta-analysis. *Journal Of Human Lactation*, 411–420.
- Dennis, C. L., & F. S. (2019). Self-Efficacy And Breastfeeding: A Systematic Review. *Journal Of Advanced Nursing*, 1843–1853.
- Donath, S. M., Amir, L. H., & Research, C. H. (2008). Maternal Obesity And Initiation And Duration Of Breastfeeding: Data From The Longitudinal Study Of Australian Children. In *Maternal And Child Nutrition* (Vol. 4).
- Ervina, A. (2018). Hubungan Paritas Dengan Asi eksklusif Pada Bayi Usia 7-12 Bulan. *Jurnal Obstetika Scienta*, 6(1), 170–178.
- Fakhidah, L. N., & Palupi, F. H. (2018). Analisis Faktor Yang Mempengaruhi Pemberian ASI Eksklusif. *Jurnal Kebidanan*, 10(2), 103–205. <http://www.jurnal.stikesseub.ac.id>
- Fauzi, F. K. (2019). Hubungan Antara Dukungan Keluarga, Status Pekerjaan Dan Paritas Ibu Dengan Perilaku Pemberian ASI Eksklusif. *Jurnal Keperawatan Muhammadiyah*, 239–243. <http://journal.um-surabaya.ac.id/index.php/JKM>
- Girard, L. C., Et Al. (2019). Parity And Breastfeeding: A Multi-Site Study. *Maternal & Child Health Journal*, 757–765.
- Guyton, A. C., & J.E., H. (2007). *Buku Ajar Fisiologi Kedokteran* (9th Ed.). EGC.
- Hasiana, S. T., Ivone, J., & Rentian, N. (2014). *Pengaruh Sectio Caesarea Terhadap Keberhasilan Asi Eksklusif The Effects Of Cesarean Delivery To The Success Of Exclusive Breastfeeding*. <http://repository.maranatha.edu/12827/>
- Hastuti, B. W., Machfudz, S., & Febriani, T. B. (2015). Hubungan Pengalaman Menyusui Dan Tingkat Pendidikan Ibu Dengan Pemberian ASI Eksklusif Di Kelurahan Barukan, Kecamatan Manisrenggo, Kabupaten Klaten. *JKKI*, 6(4), 179–187.
- Haugen, M., Et Al. (2018). Prepregnancy Body Mass Index And Breastfeeding: A Cohort Study. *BMC Pregnancy And Childbirth*, p. 15.
- Kent, J. C., Et Al. (2020). The Role Of Maternal BMI And Parity In Breastfeeding Success: A Prospective Cohort Study. *American Journal Of Clinical Nutrition*, pp. 936–945.
- Li, R., Et Al. (2021). Parity And Breastfeeding Duration In The U.S.: Findings From The Infant Feeding Practices Study II. *Pediatric Research*, pp. 123–130.
- Mabud, N., Mandang, J., & Mamuaya, T. (2014). Hubungan Pengetahuan, Pendidikan, Paritas Dengan Pemberian ASI Eksklusif Di Puskesmas Bahu Kecamatan Malalayang Kota Manado. *Jurnal Ilmiah Bidan*, 2(2), 51–56.
- Marshall, N. E., Lau, B., Purnell, J. Q., & Thornburg, K. L. (2019). Impact Of Maternal Obesity And Breastfeeding Intention On Lactation Intensity And Duration. *Maternal And Child Nutrition*, 15(2). <https://doi.org/10.1111/Mcn.12732>

- Morales, F. F., Et Al. (2021). The Combined Effect Of BMI And Parity On Breastfeeding Outcomes: A Longitudinal Study. *BMC Pregnancy And Childbirth*, p. 218.
- Ngo, L. T. H., Chou, H. F., Gau, M. L., & Liu, C. Y. (2019). Breastfeeding Self-Efficacy And Related Factors In Postpartum Vietnamese Women. *Midwifery*, pp. 70, 84–91. <https://doi.org/10.1016/j.midw.2018.12.014>
- Oddy, W. H., Li, J., Landsborough, L., Kendall, G. E., Henderson, S., & Downie, J. (2006). The Association Of Maternal Overweight And Obesity With Breastfeeding Duration. *Journal Of Pediatrics*, 149(2), 185–191. <https://doi.org/10.1016/j.jpeds.2006.04.005>
- O’Sullivan, E. J., Perrine, C. G., & Rasmussen, K. M. (2015). Early Breastfeeding Problems Mediate The Negative Association Between Maternal Obesity And Exclusive Breastfeeding At 1 And 2 Months Postpartum 1–3. *Journal Of Nutrition*, 145(10), 2369–2378. <https://doi.org/10.3945/jn.115.214619>
- P. T.-W. Wang. (2020). Railway And Road Infrastructure In The Belt And Road Initiative Countries: Estimating The Impact Of Transport Infrastructure On Economic Growth. *Transportation Research Part A: Policy And Practice*, 288--307.
- Preusting, I., Brumley, J., Odibo, L., Spatz, D. L., & Louis, J. M. (2017). Obesity As A Predictor Of Delayed Lactogenesis II. *Journal Of Human Lactation*, 33(4), 684–691. <https://doi.org/10.1177/0890334417727716>
- Rasmussen, K. M. , & K. C. L. (2018). Obesity And Breastfeeding: Causal Pathways And Implications For Future Research. *Advances In Nutrition*, 525–536.
- Rollins, N. C., Et Al. (2016). Why Invest, And What It Will Take To Improve Breastfeeding Practices? *The Lancet*, pp. 491–504.
- Torres, L. H. , Et Al. (2021). Parity And Breastfeeding Duration: A Meta-Analysis." *Journal Of Pediatric Gastroenterology And Nutrition*. *Journal Of Pediatric Gastroenterology And Nutrition*, 45–51.
- Ulfah, H. R., & Nugroho, F. S. (2020). Hubungan Usia, Pekerjaan Dan Pendidikan Ibu Dengan Pemberian Asi Eksklusif. *Jurnal Ilmiah Keperawatan*, 8(1). <https://doi.org/10.52236/Ih.V8i1.171>
- Ulina, R., & Doveriyanti, R. (2011). *Hubungan Pendidikan." Pengetahuan, Pekerjaan, Dan Dukungan Keluarga Dengan Pemberian ASI Eksklusif Di Puskesmas Sawah Lebar Kota Bengkulu Tahun 2011*.
- Untari, J. (2017). Hubungan Antara Karakteristik Ibu Dengan Pemberian Asi Eksklusif Di Wilayah Kerja Puskesmas Minggir Kabupaten Sleman The Correlation Between Mother’s Characteristic With Exclusive Breastfeeding In The Working Area Of Puskesmas Minggir District Sleman. *Jurnal Formil (Forum Ilmiah) Kesmas Respati*, 2(1).
- Wawan, A., & Dewi, M. (2011). *Teori Dan Pengukuran Pengetahuan, Sikap, Dan Perilaku Manusia*. Nuha Medika.
- WHO. (2020). Global Breastfeeding Scorecard. *World Health Organization*.
- Yuliawati, R., Kurniasari, L., & Maryam, S. (2018). Hubungan Antara Pendidikan Dan Dukungan Keluarga Dengan Kegagalan Asi Eksklusif. *Jhes (Journal Of Health Studies)*, 2(2), 73–80. <https://doi.org/10.31101/jhes.390>
- Zhu, Y., Et Al. (2021). The Effect Of Maternal BMI On Breastfeeding Practices: A Prospective Cohort Study. *Public Health Nutrition*, pp. 2185–2192.